



# VOQUEZNA<sup>®</sup>

(vonoprazan) tablets <sup>10mg</sup><sub>20mg</sub>

**For healing and heartburn relief in adults with Erosive Esophagitis and heartburn relief in adults with Non-Erosive Gastroesophageal Reflux Disease (GERD)**

**VOQUEZNA CAN KICK SOME ACID**

### **THE FIRST AND ONLY**

FDA-approved treatment of its kind, a potassium-competitive acid blocker (PCAB).

**Eligible patients may pay as little as \$25\* with the VOQUEZNA Savings Card.**


\*Patients must have commercial insurance to be eligible. Other eligibility rules apply. See [voquezna.com/savings/#terms](https://voquezna.com/savings/#terms) for complete Terms and Conditions.

## **IMPORTANT SAFETY INFORMATION**

**Do not take VOQUEZNA if you:**

- are allergic to vonoprazan or any of the other ingredients in VOQUEZNA. Allergic reaction symptoms may include trouble breathing, rash, itching, and swelling of your face, lips, tongue or throat.
- are taking a medicine that contains rilpivirine (EDURANT, COMPLERA, JULUCA, ODEFSEY) used to treat HIV-1 (Human Immunodeficiency Virus).

**Please see additional Important Safety Information throughout, as well as [Patient Information](#) and full [Prescribing Information](#) for VOQUEZNA.**



**VOQUEZNA CAN KICK  
SOME ACID**

## INDICATION

### What is VOQUEZNA?

- VOQUEZNA® (vonoprazan) is a prescription medicine used in adults:
  - for 8 weeks to heal acid-related damage to the lining of the esophagus (called Erosive Esophagitis) and for relief of heartburn related to Erosive Esophagitis.

Please see additional Important Safety Information throughout, as well as [Patient Information](#) and full [Prescribing Information](#) for VOQUEZNA.

## TABLE OF CONTENTS

**4** [VOQUEZNA WORKS DIFFERENTLY](#)

**8** [HEALING](#)

**10** [HEARTBURN RELIEF](#)

**16** [HOW TO TAKE VOQUEZNA](#)

**18** [HELPFUL TIPS](#)

**20** [SAFETY INFORMATION](#)

**22** [ABOUT GERD](#)

**30** [SAVINGS AND SUPPORT](#)



**VOQUEZNA**<sup>®</sup>  
(vonoprazan) tablets <sup>10mg</sup>  
<sup>20mg</sup>



## VOQUEZNA WORKS DIFFERENTLY

VOQUEZNA (vonoprazan) is the first and only FDA-approved treatment of its kind. Not all treatments work for everyone. If you're having ongoing symptoms, talk with your doctor about your options.



**With or  
without** food

VOQUEZNA is a type of medication called a potassium-competitive acid blocker (PCAB). Unlike most proton pump inhibitors (PPIs), PCABs can be taken with or without food.

### INDICATION (CONT'D)

#### What is VOQUEZNA?

- VOQUEZNA® (vonoprazan) is a prescription medicine used in adults:
  - for up to 6 months to maintain healing of Erosive Esophagitis and for relief of heartburn related to Erosive Esophagitis.

### VOQUEZNA in action



After taking VOQUEZNA, the medication travels to the stomach where it binds to the proton pumps.



Here, VOQUEZNA blocks potassium from binding to the pumps. This stops the proton pumps from working.



When the proton pumps can't work, they can no longer produce acid. VOQUEZNA can help suppress acid for 24 hours.

### INDICATION (CONT'D)

#### What is VOQUEZNA?

- VOQUEZNA® (vonoprazan) is a prescription medicine used in adults:
  - for 4 weeks for relief of heartburn related to gastroesophageal reflux disease (GERD).

It is not known if VOQUEZNA is safe and effective in children.

**Please see additional Important Safety Information throughout, as well as [Patient Information](#) and full [Prescribing Information](#) for VOQUEZNA.**



**VOQUEZNA®**  
(vonoprazan) tablets 10mg  
20mg



## VOQUEZNA starts to kick some acid fast

In a clinical trial of healthy adults, VOQUEZNA (vonoprazan) has been shown to reduce acid within 2-3 hours and maintain acid suppression over 24 hours.

 **Acid reduction in**   
**2-3 HOURS**

### IMPORTANT SAFETY INFORMATION (CONT'D)

Before taking VOQUEZNA, tell your healthcare provider about all your medical conditions, including if you:

- have low magnesium, calcium, or potassium in your blood, or you are taking a medicine to increase urine (diuretic).
- have kidney or liver problems.



## VOQUEZNA healed my Erosive GERD.\*

KEVIN, VOQUEZNA Clinical Trial Patient with Erosive GERD



\*Individual results may vary.  
The patient was compensated by Phathom for his time sharing his individual experience in the trial.

[Check out more of Kevin's story >](#)


Please see additional Important Safety Information throughout, as well as [Patient Information](#) and full [Prescribing Information](#) for VOQUEZNA.

 **VOQUEZNA**<sup>®</sup>  
(vonoprazan) tablets <sup>10mg</sup><sub>20mg</sub>



## VOQUEZNA CAN KICK SOME ACID AND HELP HEAL AND MAINTAIN HEALING OF EROSIVE GERD

VOQUEZNA (vonoprazan) is shown to help most adults with Erosive GERD (also known as Erosive Esophagitis or EE) achieve complete healing that can last.



Representation of an **esophageal erosion**

### IMPORTANT SAFETY INFORMATION (CONT'D)

Before taking VOQUEZNA, tell your healthcare provider about all your medical conditions, including if you:

- are pregnant, think you may be pregnant, or plan to become pregnant. It is not known if VOQUEZNA will harm your unborn baby.

## Proven healing of Erosive GERD

VOQUEZNA was evaluated in a clinical trial of adults with endoscopically confirmed EE. One group took VOQUEZNA 20 mg once daily for 2 to 8 weeks during the healing phase. **93% of adults taking VOQUEZNA for EE experienced healing by 2 months** (compared to 85% taking another treatment\*).

Some adults who were healed by VOQUEZNA or another treatment during the healing phase then took VOQUEZNA 10 mg once daily for 6 months during the maintenance phase. **79% of those stayed healed for 6 months on VOQUEZNA maintenance treatment** (compared to 72% taking another treatment\*).



**93% OF ADULTS**

TAKING VOQUEZNA ACHIEVED **EE HEALING BY 2 MONTHS** VS 85% TAKING ANOTHER TREATMENT\*

AND OF THOSE HEALED



**79% OF ADULTS**

TAKING VOQUEZNA **STAYED HEALED FOR 6 MONTHS** VS 72% TAKING ANOTHER TREATMENT\*

\*lansoprazole

### IMPORTANT SAFETY INFORMATION (CONT'D)

- **Pregnancy Exposure Registry:** There is a pregnancy registry for women who take VOQUEZNA during pregnancy. The purpose of this registry is to collect information about the the health of you and your baby.

Please see additional Important Safety Information throughout, as well as [Patient Information](#) and full [Prescribing Information](#) for VOQUEZNA.



**VOQUEZNA**<sup>®</sup>  
(vonoprazan) tablets <sup>10mg</sup>  
<sup>20mg</sup>



## HELPING PATIENTS ACHIEVE HEARTBURN-FREE DAYS

VOQUEZNA (vonoprazan) is proven to help adults with Erosive\* or Non-Erosive GERD achieve heartburn-free days.

\*Erosive GERD is also known as Erosive Esophagitis or EE.

### IMPORTANT SAFETY INFORMATION (CONT'D)

**Before taking VOQUEZNA, tell your healthcare provider about all your medical conditions, including if you:**

- are breastfeeding or plan to breastfeed. You and your healthcare provider should decide if you will take VOQUEZNA while breastfeeding.

**Tell your healthcare provider about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Keep a list of them to show your healthcare provider and pharmacist when you get a new medicine.



**PROVEN TO HELP RELIEVE HEARTBURN FOR GERD**

### IMPORTANT SAFETY INFORMATION (CONT'D)

VOQUEZNA may affect how other medicines work, and other medicines may affect how VOQUEZNA works. Especially tell your healthcare provider if you take medicine that contains rilpivirine (EDURANT, COMPLERA, JULUCA, ODEFSEY).

**Please see additional Important Safety Information throughout, as well as [Patient Information](#) and full [Prescribing Information](#) for VOQUEZNA.**

 **VOQUEZNA<sup>®</sup>**  
(vonoprazan) tablets 10mg  
20mg

## Heartburn relief in adults with Non-Erosive GERD

In a clinical trial of adults with Non-Erosive GERD (Gastroesophageal Reflux Disease) taking VOQUEZNA (vonoprazan), subjects were asked to keep track of the days they experienced **heartburn and the severity of heartburn symptoms**. Diaries were checked at the end of treatment to identify how many days were reported to be heartburn free.



**45% of days spent on VOQUEZNA were heartburn free for 24 hours** (compared to 28% of days on placebo).

### IMPORTANT SAFETY INFORMATION (CONT'D)

What are the possible side effects of VOQUEZNA?

VOQUEZNA may cause serious side effects including:

- **A type of kidney problem (acute tubulointerstitial nephritis):** Some people who take VOQUEZNA may develop a kidney problem called acute tubulointerstitial nephritis. Call your healthcare provider right away if you have a decrease in the amount that you urinate or if you notice blood in your urine.
- **Diarrhea caused by an infection (*Clostridioides difficile*) in your intestines:** Call your healthcare provider right away if you have watery stools, stomach pain, and fever that does not go away.
- **Bone fractures (hip, wrist, or spine):** Bone fractures in the hip, wrist, or spine may happen in people who take multiple daily doses of another type of medicine that reduces acid in your stomach



**I'm happy to report that, thanks to VOQUEZNA, my heartburn no longer takes the center stage in my life.\***

**CARSON**, VOQUEZNA Clinical Trial Patient with Non-Erosive GERD



\*Individual results may vary.  
The patient was compensated by Phathom for his time sharing his individual experience in the trial.

### IMPORTANT SAFETY INFORMATION (CONT'D)

known as proton pump inhibitors (PPI medicines) for a long period of time (a year or longer). Tell your healthcare provider if you have a bone fracture, especially in the hip, wrist, or spine.

Please see additional Important Safety Information throughout, as well as [Patient Information](#) and full [Prescribing Information](#) for VOQUEZNA.



**VOQUEZNA**<sup>®</sup>  
(vonoprazan) tablets 10mg  
20mg

## Heartburn relief in adults with Erosive GERD

In a clinical trial of adults with Erosive GERD (also known as Erosive Esophagitis or EE) taking VOQUEZNA (vonoprazan), subjects were asked to keep track of the days they experienced **heartburn and the severity of heartburn symptoms**.

Diaries were checked at the end of treatment to identify how many days were reported to be heartburn free.



During the 2-month healing phase, **67% of days spent on VOQUEZNA were heartburn free for 24 hours** (compared to 64% of days on another treatment\*). Over the 6-month maintenance phase, **81% of days spent on VOQUEZNA were heartburn free for 24 hours** (compared to 79% of days on another treatment\*).

\*lansoprazole

### IMPORTANT SAFETY INFORMATION (CONT'D)

#### VOQUEZNA may cause serious side effects including:

- **Severe skin reactions:** VOQUEZNA can cause rare, but severe skin reactions that may affect any part of your body. These serious skin reactions may need to be treated in a hospital and may be life threatening:
  - Skin rash which may have blistering, peeling, or bleeding on any part of your skin.
  - You may also have fever, chills, body aches, shortness of breath, or enlarged lymph nodes.

If you experience any of these symptoms, stop taking VOQUEZNA and call your healthcare provider right away. These symptoms may be the first sign of a severe skin reaction.

### IMPORTANT SAFETY INFORMATION (CONT'D)

#### VOQUEZNA may cause serious side effects including:

- **Low vitamin B-12 levels:** VOQUEZNA lowers the amount of acid in your stomach. Stomach acid is needed to absorb Vitamin B12 properly. Tell your healthcare provider if you have symptoms of low vitamin B12 levels, including irregular heartbeat, shortness of breath, lightheadedness, tingling or numbness in the arms or legs, muscle weakness, pale skin, feeling tired, or mood changes. Talk with your healthcare provider about the risk of low vitamin B12 levels if you have been on VOQUEZNA for a long time.

**Please see additional Important Safety Information throughout, as well as [Patient Information](#) and full [Prescribing Information](#) for VOQUEZNA.**



**VOQUEZNA<sup>®</sup>**  
(vonoprazan) tablets <sup>10mg</sup>  
<sup>20mg</sup>



## HOW TO TAKE VOQUEZNA



VOQUEZNA (vonoprazan) is a once-daily pill for adults with heartburn related to both Erosive or Non-Erosive GERD.

**It can be taken with or without food.**

Your dosing schedule may look something like what is outlined on the next page.

### IMPORTANT SAFETY INFORMATION (CONT'D)

**VOQUEZNA may cause serious side effects including:**

- **Low magnesium levels in the body** can happen in people who take VOQUEZNA. Tell your healthcare provider right away if you have symptoms of low magnesium levels, including seizures, dizziness, irregular heartbeat, jitteriness, muscle aches or weakness, or spasms of the hands, feet, or voice.

## Dosing plan

### RELIEF OF HEARTBURN RELATED TO NON-EROSIVE GERD



One 10 mg pill  
Once daily  
For 4 weeks

### HEALING OF EROSIVE GERD AND RELIEF OF RELATED HEARTBURN



One 20 mg pill  
Once daily  
For 8 weeks

### MAINTENANCE OF HEALED EROSIVE GERD AND RELIEF OF RELATED HEARTBURN



One 10 mg pill  
Once daily  
Up to 6 months

Pills shown not actual size.

Talk to your doctor to determine a management plan that's right for you.

### IMPORTANT SAFETY INFORMATION (CONT'D)

- **Stomach growths (fundic gland polyps):** A certain type of stomach growth called fundic gland polyps may happen in people who take another type of medicine that reduces acid in your stomach known as proton pump inhibitors (PPI medicines) for a long time. Talk with your healthcare provider about the possibility of fundic gland polyps if you have been on VOQUEZNA for a long time.

**Please see additional Important Safety Information throughout, as well as [Patient Information](#) and full [Prescribing Information](#) for VOQUEZNA.**



# HELPFUL TIPS WHEN TAKING VOQUEZNA



## Flexibility to take with or without food

VOQUEZNA (vonoprazan) is a once-daily pill that can be taken with or without food, unlike most proton pump inhibitors (PPIs), which means you do not have to take it around mealtimes. Remember, you should always take VOQUEZNA as your doctor prescribes.

## Take it whole

Chewing or crushing the tablet may cause a bitter taste.

Pill shown not actual size or color.



## IMPORTANT SAFETY INFORMATION (CONT'D)

The most common side effects of VOQUEZNA for treatment of Erosive Esophagitis or relief of heartburn related to gastroesophageal reflux disease include:

- stomach inflammation
- indigestion
- diarrhea
- constipation
- stomach bloating
- high blood pressure
- stomach pain
- urinary tract infection
- nausea



## TIP:

Try taking VOQUEZNA right after an activity you do once a day. It can help you incorporate it into your routine.

## Miss a dose?



If you miss a dose, take the next as soon as possible—unless 12 hours have passed. If more than 12 hours have passed, skip the missed dose and take your next dose on time.

## IMPORTANT SAFETY INFORMATION (CONT'D)

These are not all the possible side effects of VOQUEZNA. For more information, ask your healthcare provider or pharmacist. Call your healthcare provider for medical advice about side effects.

You are encouraged to report suspected adverse reactions by contacting Phathom Pharmaceuticals at **1-888-775-PHAT (7428)** or FDA at **1-800-FDA-1088** or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

Please see additional Important Safety Information throughout, as well as [Patient Information](#) and full [Prescribing Information](#) for VOQUEZNA.





## SAFETY AND SIDE EFFECTS

VOQUEZNA (vonoprazan) was evaluated in clinical trials of adults with either Non-Erosive Gastroesophageal Reflux Disease (GERD) or Erosive GERD (also known as Erosive Esophagitis or EE).

The most common side effects listed on the next page are not the full list of possible side effects. To see the full list of possible side effects, please see full [Prescribing Information](#).

### Non-Erosive GERD (4 weeks)

The most common side effects occurring in adults taking VOQUEZNA 10 mg were:

Side effects	VOQUEZNA 10 mg 259 adults
Stomach pain	2%
Constipation	2%
Diarrhea	2%
Nausea	2%
Urinary tract infection	2%

### Erosive GERD healing phase (up to 2 months)

The most common side effects occurring in adults taking VOQUEZNA 20 mg were:

Side effects	VOQUEZNA 20 mg 514 adults
Stomach inflammation	3%
Diarrhea	2%
Stomach bloating	2%
Stomach pain	2%
Nausea	2%

### Erosive GERD maintenance phase (6 months)

The most common side effects occurring in adults taking VOQUEZNA 10 mg were:

Side effects	VOQUEZNA 10 mg 296 adults
Stomach inflammation	6%
Stomach pain	4%
Indigestion	4%
High blood pressure	3%
Urinary tract infection	3%

Please see additional Important Safety Information throughout, as well as [Patient Information](#) and full [Prescribing Information](#) for VOQUEZNA.





## WHAT TO KNOW ABOUT GERD

Gastroesophageal Reflux Disease (GERD) is a chronic disease that happens when stomach acid flows into the esophagus and causes irritation.

Two main types of GERD are:

- Non-Erosive GERD
- Erosive GERD (also known as Erosive Esophagitis or EE)

## Who does GERD affect?

GERD is one of the most common digestive diseases, affecting approximately 65 million people in the United States.



### Erosive GERD vs Non-Erosive GERD

Erosive GERD is a type of GERD in which acid-related damage to the esophagus occurs. Non-Erosive GERD does not involve damage to the esophagus and is more common. Heartburn is common in both conditions.

**~2/3 OF PEOPLE**

LIVING WITH GERD HAVE NON-EROSIVE GERD

### Heartburn that persists

Many people with Erosive GERD continue to feel acid in their esophagus despite taking medicines such as antacids, histamine-2 receptor antagonists (H2s), or proton pump inhibitors (PPIs).



**58% OF PEOPLE**

SUFFERING FROM EROSIIVE GERD STILL EXPERIENCE HEARTBURN WHILE ON THEIR TREATMENT

## Factors that may increase your chances of developing Gastroesophageal Reflux Disease (GERD) include:

- Hiatal hernia
- Obesity
- Lower esophageal sphincter doesn't close correctly
- Delayed stomach emptying

## What symptoms are related to GERD?



**The most common symptoms are heartburn and regurgitation. Other possible symptoms include:**

- Difficult or painful swallowing
- Chronic cough
- Chest pain

## How is GERD diagnosed?

**To diagnose GERD**, your doctor will ask you a few questions and may perform a physical exam.

**To diagnose Erosive GERD**, an endoscopy is needed. An endoscopy is a procedure in which a thin tube equipped with a tiny camera is lowered down your throat.



## The history of GERD treatments

### 1928

**The first antacid was approved for use**

Antacids neutralize the acid in your stomach to help relieve heartburn. They are typically offered over the counter and are used for symptom relief.

### 1977

**The first histamine-2 receptor antagonist (H2) was approved for use**

H2s reduce the amount of acid produced in the stomach by blocking one of the first steps in acid production. An H2 treats heartburn symptoms related to GERD.

### 1989

**The first proton pump inhibitor (PPI) was approved for use**

PPIs are a common treatment for GERD. They treat GERD by blocking the final step of acid production. They are activated by stomach acid, and most PPIs must be taken 30-60 minutes prior to meals.

### 2023

### 2024

**VOQUEZNA® (vonoprazan) was approved for use as the first potassium-competitive acid blocker**

VOQUEZNA was approved as the first potassium-competitive acid blocker (PCAB) to treat Erosive Esophagitis and its symptoms in 2023 and later to relieve heartburn caused by Non-Erosive GERD in 2024. PCABs work by blocking the final step of acid production—they do not need to be activated by stomach acid and can be taken with or without food.

## Talking to your doctor

When it comes to GERD (Gastroesophageal Reflux Disease), not every treatment works for everyone. If you've been wondering if a different treatment option is right for you, it may be time to talk to your doctor.



### ✓ **Be timely.**

The longer you delay talking to your doctor about troublesome symptoms, the longer you could be delaying healing of your Erosive GERD damage or related heartburn relief.

### ✓ **Be honest.**

The best way for your doctor to find a treatment that works for you is by collecting all the facts. Don't be shy with your doctor. They're your best ally.

### ✓ **Be descriptive.**

No detail is too small. What time of day do your symptoms occur? How long, often, and bothersome are your symptoms? Tell your doctor these details so they can identify the best treatment for you.

## INDICATION AND IMPORTANT SAFETY INFORMATION

### What is VOQUEZNA?

- VOQUEZNA® (vonoprazan) is a prescription medicine used in adults:
  - for 8 weeks to heal acid-related damage to the lining of the esophagus (called Erosive Esophagitis) and for relief of heartburn related to Erosive Esophagitis.
  - for up to 6 months to maintain healing of Erosive Esophagitis and for relief of heartburn related to Erosive Esophagitis.
  - for 4 weeks for relief of heartburn related to gastroesophageal reflux disease (GERD).

It is not known if VOQUEZNA is safe and effective in children.

### Do not take VOQUEZNA if you:

- are allergic to vonoprazan or any of the other ingredients in VOQUEZNA. Allergic reaction symptoms may include trouble breathing, rash, itching, and swelling of your face, lips, tongue or throat.
- are taking a medicine that contains rilpivirine (EDURANT, COMPLERA, JULUCA, ODEFSEY) used to treat HIV-1 (Human Immunodeficiency Virus).

### Before taking VOQUEZNA, tell your healthcare provider about all your medical conditions, including if you:

- have low magnesium, calcium, or potassium in your blood, or you are taking a medicine to increase urine (diuretic).
- have kidney or liver problems.
- are pregnant, think you may be pregnant, or plan to become pregnant. It is not known if VOQUEZNA will harm your unborn baby.
  - **Pregnancy Exposure Registry:** There is a pregnancy registry for women who take VOQUEZNA during pregnancy. The purpose of this registry is to collect information about the health of you and your baby. If you are pregnant or become pregnant during treatment with VOQUEZNA, talk to your healthcare provider about how you can join this pregnancy registry or you can enroll in the registry by calling 1-866-609-1612 or visiting <https://voqueznapregnancyregistry.com/>.



## IMPORTANT SAFETY INFORMATION (CONT'D)

### Before taking VOQUEZNA, tell your healthcare provider about all your medical conditions, including if you:

- are breastfeeding or plan to breastfeed. You and your healthcare provider should decide if you will take VOQUEZNA while breastfeeding.

**Tell your healthcare provider about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Keep a list of them to show your healthcare provider and pharmacist when you get a new medicine.

VOQUEZNA may affect how other medicines work, and other medicines may affect how VOQUEZNA works. Especially tell your healthcare provider if you take medicine that contains rilpivirine (EDURANT, COMPLERA, JULUCA, ODEFSEY).

### What are the possible side effects of VOQUEZNA?

#### VOQUEZNA may cause serious side effects including:

- **A type of kidney problem (acute tubulointerstitial nephritis):** Some people who take VOQUEZNA may develop a kidney problem called acute tubulointerstitial nephritis. Call your healthcare provider right away if you have a decrease in the amount that you urinate or if you notice blood in your urine.
- **Diarrhea caused by an infection (*Clostridioides difficile*) in your intestines:** Call your healthcare provider right away if you have watery stools, stomach pain, and fever that does not go away.
- **Bone fractures (hip, wrist, or spine):** Bone fractures in the hip, wrist, or spine may happen in people who take multiple daily doses of another type of medicine that reduces acid in your stomach known as proton pump inhibitors (PPI medicines) for a long period of time (a year or longer). Tell your healthcare provider if you have a bone fracture, especially in the hip, wrist, or spine.
- **Severe skin reactions:** VOQUEZNA can cause rare, but severe skin reactions that may affect any part of your body. These serious skin reactions may need to be treated in a hospital and may be life threatening:
  - Skin rash which may have blistering, peeling, or bleeding on any part of your skin.
  - You may also have fever, chills, body aches, shortness of breath, or enlarged lymph nodes.

If you experience any of these symptoms, stop taking VOQUEZNA and call your healthcare provider right away. These symptoms may be the first sign of a severe skin reaction.

- **Low vitamin B-12 levels:** VOQUEZNA lowers the amount of acid in your stomach. Stomach acid is needed to absorb Vitamin B12 properly. Tell your healthcare provider if you have symptoms of low vitamin B12 levels, including irregular heartbeat, shortness of breath, lightheadedness, tingling or numbness in the arms or legs, muscle weakness, pale skin, feeling tired, or mood changes. Talk with your healthcare provider about the risk of low vitamin B12 levels if you have been on VOQUEZNA for a long time.
- **Low magnesium levels in the body** can happen in people who take VOQUEZNA. Tell your healthcare provider right away if you have symptoms of low magnesium levels, including seizures, dizziness, irregular heartbeat, jitteriness, muscle aches or weakness, or spasms of the hands, feet, or voice.
- **Stomach growths (fundic gland polyps):** A certain type of stomach growth called fundic gland polyps may happen in people who take another type of medicine that reduces acid in your stomach known as proton pump inhibitors (PPI medicines) for a long time. Talk with your healthcare provider about the possibility of fundic gland polyps if you have been on VOQUEZNA for a long time.

#### The most common side effects of VOQUEZNA for treatment of Erosive Esophagitis or relief of heartburn related to gastroesophageal reflux disease include:

- stomach inflammation
- indigestion
- diarrhea
- constipation
- stomach bloating
- high blood pressure
- stomach pain
- urinary tract infection
- nausea

These are not all the possible side effects of VOQUEZNA. For more information, ask your healthcare provider or pharmacist. Call your healthcare provider for medical advice about side effects.

**You are encouraged to report suspected adverse reactions by contacting Phathom Pharmaceuticals at 1-888-775-PHAT (7428) or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

**Please see additional Important Safety Information throughout, as well as [Patient Information](#) and full [Prescribing Information](#) for VOQUEZNA.**





## YOU MAY SAVE ON YOUR VOQUEZNA PRESCRIPTION

With the VOQUEZNA Savings Card, eligible commercially insured patients may **pay as little as \$25\*** for their prescription.



You can download a Savings Card and sign up for additional support here >

**BLINKRx**

Our online  
pharmacy partner

BlinkRx helps identify potential savings available to you. Phathom Pharmaceuticals and BlinkRx work together to help simplify the process for receiving VOQUEZNA (vonoprazan).

If your doctor sends your prescription to BlinkRx, BlinkRx will contact you to begin setting up free home delivery.



### **FREE home delivery**

A pharmacy partner will deliver your prescription FREE to your door.



### **Your lowest out-of-pocket cost**

BlinkRx will identify your medication cost and apply eligible savings offers.

To see full Terms & Conditions please go to [voquezna.com/savings/#terms](https://voquezna.com/savings/#terms)

BlinkRx is NOT insurance. By sending your prescription to BlinkRx, you authorize BlinkRx ("Blink") to send text message notifications related to your prescriptions. Message and data rates may apply. Message frequency varies. Standard SMS messaging may not be fully secure. Providing consent is not a condition of purchasing goods or services through Blink.

You can reply to any text with HELP for help or STOP to stop and revoke your consent. Visit Blink's HIPAA Privacy Notice and Terms of Use at [www.blinkrx.com/legal](https://www.blinkrx.com/legal).



# A DIFFERENT TREATMENT OPTION

- ✓ **The first and only FDA-approved** treatment of its kind, a potassium-competitive acid blocker (PCAB).
- ✓ **Can help relieve heartburn** related to Non-Erosive GERD (Gastroesophageal Reflux Disease) or Erosive GERD (also known as Erosive Esophagitis or EE).
- ✓ **Can help heal and maintain healing** of Erosive GERD for most adults.

## The most common side effects

for adults with Non-Erosive GERD or EE were stomach inflammation, diarrhea, stomach bloating, stomach pain, nausea, indigestion, constipation, high blood pressure, and urinary tract infection. Other serious side effects have occurred. Please see [Patient Information](#) for other possible side effects.

## Eligible patients may pay as little as \$25\* with the VOQUEZNA Savings Card

\*Patients must have commercial insurance to be eligible. Other eligibility rules apply. See [voquezna.com/savings/#terms](https://voquezna.com/savings/#terms) for complete Terms and Conditions.

## IMPORTANT SAFETY INFORMATION

### Do not take VOQUEZNA if you:

- are allergic to vonoprazan or any of the other ingredients in VOQUEZNA. Allergic reaction symptoms may include trouble breathing, rash, itching, and swelling of your face, lips, tongue or throat.
- are taking a medicine that contains rilpivirine (EDURANT, COMPLERA, JULUCA, ODEFSEY) used to treat HIV-1 (Human Immunodeficiency Virus).

Please see additional Important Safety Information throughout, as well as [Patient Information](#) and full [Prescribing Information](#) for VOQUEZNA.